

DUST EVOLUTION IN PROTOSTELLAR COLLAPSE

Maxime Lombart (maxime.lombart@cea.fr)
postdoc in ERC project PEBBLES/CEA Paris-Saclay

Wada et al. 2018

With: U. Lebreuilly, A. Maury

Collaborators: C.-E. Bréhier, M.
Hutchison, Y.-N. Lee, G. Laibe,
D. J. Price




RAMSES SNO days, 2025/11/25, Paris



STAR FORMATION, DISK EVOLUTION AND PLANET FORMATION


The first seeds for planets may form in less than a few million years




Dust growth must be an efficient process !

3D SIMULATIONS WITH DUST COAGULATION/FRAGMENTATION


discs



Vericel et al., 2021




protostellar
collapses



Lebreuilly et al., 2023



Tsukamoto et al., 2021



gas and dust dynamic in 1D

single dust size approximation

dust size distribution

gas and dust dynamic in 3D



3D SIMULATIONS WITH DUST COAGULATION


Marian Smoluchowski (1872-1917)



m, m' : masses

n : number density
of particles per
mass unit

K : collision kernel



$$\frac{\partial n(m, t)}{\partial t} = \underbrace{\frac{1}{2} \int_0^m K(m', m - m') n(m', t) n(m - m', t) dm'}_{\text{gain}} - \underbrace{n(m, t) \int_0^\infty K(m, m') n(m', t) dm'}_{\text{loss}}$$

Non-linear integro-differential equation !!! 🤯

In astrophysics: $K(m_1, m_2) = \sigma(m_1, m_2) \Delta v(m_1, m_2) \rightarrow$ need numerical solutions



Requirements

- ◇ 3 orders of magnitude in size ($1\mu\text{m} - 1\text{mm}$) $\Leftrightarrow$ 9 orders in mass
- ◇ mass range discretization $\rightarrow$ 20 mass bins (low numerical cost)

HIGH-ORDER NUMERICAL SCHEME

Discontinuous Galerkin scheme


conservative form (Tanaka et al., 1996)

Liu et al., 2019
Lombart &
Laibe, 2021

$$\begin{cases} \frac{\partial g(m, t)}{\partial t} + \frac{\partial F_{\text{coag}}[g](m, t)}{\partial m} = 0 \\ F_{\text{coag}}[g](m, t) = \int_0^m \int_{m-m_1}^{\infty} \frac{K(m_1, m_2)}{m_2} g(m_1, t) g(m_2, t) dm_2 dm_1 \end{cases}$$




COAgulation with
Lightning fast
Algorithm



→ high accuracy
→ few number of mass bins

BENCHMARK COALA

Lombart & Laibe (2021), Laibe & Lombart (2022)



Requirements

- ✓ 9 orders of magnitude in mass
- ✓ 10-20 dust size bins

3D simulations with
polydisperse coagulation



DUST GROWTH IN PROTOSTELLAR COLLAPSE

RAMSES MHD model of B335-like
protostar
gas & dust (40 bins)

Implementation:

- dust growth as source term of continuity equation

$$\frac{\partial \rho_k}{\partial t} + \nabla \cdot [\rho_k \mathbf{v}_k] = \left. \frac{\partial \rho_k}{\partial t} \right|_{\text{growth}}$$


$$\rho_k = \int_{I_k} g(m, t) dm$$

- ordinary operator splitting

$$\rho_k^* = \rho_k(t_n) - \Delta t \left(\nabla \cdot [\rho_k \mathbf{v}_k] \right) \rightarrow \text{RAMSES}$$



$$\rho_k(t_{n+1}) = \rho_k^* + \Delta t \left. \frac{\partial \rho_k}{\partial t} \right|_{\text{growth}} \rightarrow \text{COALA}$$

Lombart et al., 2025 (minor revisions)



DUST GROWTH IN PROTOSTELLAR COLLAPSE

Lombart et al., 2025 (minor revisions)





Mass of grains in the envelope
(in % of total dust mass)

Initially (MRN distribution):

- $\sim 0.25 \mu\text{m} \rightarrow 20\%$
- $\sim 5 \mu\text{m} \rightarrow 0\%$

After 100 kyrs:

- $\sim 0.25 \mu\text{m} \rightarrow 13\%$
- $\sim 5 \mu\text{m} \rightarrow 1\%$



DUST FRAGMENTATION MODEL

Destructive fragmentation



Mass transfer



Mass transfer + cratering



$$\frac{\partial n(m, t)}{\partial t} = \frac{1}{2} \int_0^\infty \int_0^\infty \underbrace{K(m_1, m_2) b(m, m_1, m_2)}_{\text{mass distribution of fragments}} n(m_1, t) n(m_2, t) dm_1 dm_2 - n(m, t) \int_0^\infty K(m, m_1) n(m_1, t) dm_1$$




Safronov, 1972; Jones et al., 1996; Suttner et al., 2001; Blum, 2006; Hirashita et al., 2009; Rafikov et al., 2020

Conservative form (Lombart et al. 2024):

$$F_{\text{frag}}[g](m, t) = \frac{1}{2} \int_0^m \int_0^\infty \int_0^\infty \mathbf{1}_{m_1+m_2 \geq m} \frac{m_f}{m_1 m_2} b(m_f, m_1, m_2) K(m_1, m_2) g(m_1, t) g(m_2, t) dm_2 dm_1 dm_f$$

$$- \int_0^m \int_{m-m_1}^\infty \frac{K(m_1, m_2)}{m_2} g(m_1, t) g(m_2, t) dm_2 dm_1$$

Bukhari Syed et al. 2017



DUST FRAGMENTATION MODEL

COALA

Benchmark for
fragmentation

Feingold et al., 1988


$$K(u, v) = 1$$

$$b(w, u, v) = \gamma^2(u + v)e^{-\gamma w}$$

3D simulations
with polydisperse
general non-linear
fragmentation




Lombart et al. 2024




DUST MODEL IMPROVEMENT

Growth and fragmentation of multi-component dust aggregates



MULTI-COMPONENT DUST COAGULATION



Need 2D Smoluchowski
coagulation equation

$$\frac{\partial n(m_s, m_i, t)}{\partial t} = \frac{1}{2} \int_0^{m_s} \int_0^{m_i} K(m_s - m'_s, m'_s, m_i - m'_i, m'_i) n(m'_s, m'_i, t) n(m_s - m'_s, m_i - m'_i, t) dm'_s dm'_i - \int_0^\infty \int_0^\infty K(m_s, m'_s, m_i, m'_i) n(m_s, m_i, t) n(m'_s, m'_i, t) dm'_s dm'_i$$

$$K(m_s - m'_s, m'_s, m_i - m'_i, m'_i) = P_{\text{stick}}(m_s, m'_s, m_i, m'_i) \sigma \Delta v$$

Sticking
probability $\rightarrow$

$$P_{\text{stick}}(m_s, m'_s, m_i, m'_i) = \epsilon_{s-s} \frac{m_{s,A}}{m_A} \frac{m_{s,B}}{m_B} + \epsilon_{s-i} \left(\frac{m_{s,A}}{m_A} \frac{m_{i,A}}{m_A} + \frac{m_{s,B}}{m_B} \frac{m_{i,B}}{m_B} \right) + \epsilon_{i-i} \frac{m_{i,A}}{m_A} \frac{m_{i,B}}{m_B}$$

Sticking efficiency
of s-s collision


Sticking efficiency
of s-i collision

Sticking efficiency
of i-i collision

Need data from
lab experiments

DUST AGGREGATION

Dust aggregate is defined by its mass m and porosity $\psi = \frac{V}{V^*}$



Need 2D Smoluchowski coagulation equation

$$\frac{\partial n(m, \psi_m, \tau)}{\partial t} = \frac{1}{2} \int_0^m \int_{\psi_{\min}}^{\psi_{\max}} \int_{\psi_{\min}}^{\psi_{\max}} K(m - m_1, \psi_{m-m_1}; m_1, \psi_{m_1}) n(m_1, \psi_{m_1}, t) n(m - m_1, \psi_{m-m_1}, t) \delta(\psi_m - \Gamma(m - m_1, \psi_{m-m_1}; m_1, \psi_{m_1})) d\psi_{m-m_1} d\psi_{m_1} dm_1$$

$$- n(m, \psi_m, t) \int_0^\infty \int_{\psi_{\min}}^{\psi_{\max}} K(m, \psi_m; m_1, \psi_{m_1}) n(m_1, \psi_{m_1}, t) d\psi_{m_1} dm_1$$

$\Gamma(m - m_1, \psi_{m-m_1}; m_1, \psi_{m_1}) \rightarrow$ determine the porosity of the resulting aggregate



Blum, 2006; Ormel et al., 2007; Okuzumi et al., 2009; Hirashita et al., 2021


FRAGMENTATION OF MULTI-COMPONENT DUST AGGREGATES

Multi-component dust:

→ how to model the composition of fragments ?

Dust aggregates:

→ how to model the mass and porosity of the fragments ?




Need data from lab experiments



Blum, 2006; Ormel et al., 2007; Okuzumi et al., 2009; Hirashita et al., 2021; Hasegawa et al., 2022



CHARACTERIZING DUST EVOLUTION DURING THE DISK-BUILDING STAGE



Carpine et al., 2025

Realistic dust
optical properties




THEMIS

The Heterogeneous dust Evolution Model for
Interstellar Solids



amorphous silicate core
+ carbon mantle

amorphous carbon
hydrogenated

Monomers
 $\sim 0.1 \mu\text{m}$



$\sim 1 \mu\text{m}$



aggregates

compact sphere

Radiative transfer
codes

Realistic synthetic
observations

TAKE HOME MESSAGES

- The **size evolution** of dust particles are modeled by **Smoluchowski-like equations**.
- **3D simulations with polydisperse dust coagulation/fragmentation**
 - 9 orders of magnitude in mass ($1\mu\text{m}$ - 1mm)
 - Few number of bins ~ 20
 - Good accuracy
- **Dust grains are predicted to grow by 2 orders of magnitude in collapsing envelopes (just gravity/turbulence, without magnetic effects yet)**
- Coagulation/fragmentation of **dust aggregates** and **multicomponent dust** can be modeled by **2D extension**.
- **PEBBLES**: realistic synthetic observations by combining simulations, radiative transfer with realistic dust optical properties

